



### FOREIGN TECHNOLOGY DIVISION



UNSTEADY DOWNWASH BEHIND A DELTA WING WITH SUPERSONIC MOTION

Ву

R. Sh. Solomonyan





Approved for public release; distribution unlimited.

78 12 26

AD-4066488

29

# UNEDITED MACHINE TRANSLATION

FTD-ID(RS)T-1842-78

2 November 1978

MICROFICHE NR: 34D-78-C-00 1493

CSP73108310

UNSTEADY DOWNWASH BEHIND A DELTA WING WITH SUPERSONIC MOTION

By: R. Sh. Solomonyan

English pages: 26

Izvestiya Akademii Nauk Armyanskoy SSR,

Mekhanika, Vol. 25, Nr. 5, 1972,

pp. 45-64

Country of Origin: USSR

This document is a machine translation.

Requester: FTD/TQTA

Approved for public release; distribution

unlimited.

THIS TRANSLATION IS A RENDITION OF THE ORIGI-NAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DI-VISION.

PREPARED BY:

TRANSLATION DIVISION FOREIGN TECHNOLOGY DIVISION WP-AFB, OHIO.

U. S. BOARD ON GEOGRAPHIC NAMES TRANSLITERATION SYSTEM

Block	Italic	Transliteration	Block	Italic	Transliteration
A a	Aa	A, a	Рр	PP	R, r
Бб	5 6	B, b	Сс	Cc	S, s
Вв	B .	V, v	Тт	T m	T, t
Гг	Γ .	G, g	Уу	уу	U, u
Дд	Дд	D, d	Фф	<b>•</b> •	F, f
Еe	E .	Ye, ye; E, e*	X x	Xx	Kh, kh
жж	Ж ж	Zh, zh	Цц	4 4	Ts, ts
3 з	3 ;	Z, z	4 4	4 4	Ch, ch
Ии	и и	I, i	Шш	Шш	Sh, sh
Йй	A a	Ү, у	Щщ	Щщ	Shch, shch
Нн	K x	K, k	Ъъ	3 .	11
ת ונ	ЛА	L, 1	Н ы	M W	У, у
in in	M M	M, m	Ьь	b .	•
Нн.	H N	N, n	Ээ	9 ,	Е, е
0 0	0 0	0, 0	Юю	10 no	Yu, yu
Пп	Пп	P, p	Яя	Яя	Ya, ya

\*ye initially, after vowels, and after ъ, ь; e elsewhere. When written as  $\ddot{e}$  in Russian, transliterate as  $y\ddot{e}$  or  $\ddot{e}$ .

### RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	sinh-1
cos tg	tan	th	cosh tanh	arc ch arc th	tanh-1
ctg	cot	cth sch	coth sech	arc cth arc sch	sech_1
cosec	csc	csch	csch	arc csch	csch-1

Russian	English
rot	curl
1g	log

Fage 45.

UNSTRADY DOWNWASH BEHIND A DELTA WING WITH SUPERSONIC MOTION.

B. Sh. Solomonyan

The question of determining the aerodynamic characteristics of the tail assembly of flight vehicle has great practical value both from the point of view of control of flight vehicles and from the point of view of aeroelasticity.

During the determination of the aerodynamic forces and tail moments, arises the question of the determination of wing downwash. This task in special cases is examined in works [5, 7], etc., but common/general/total setting is given by N: N. Kislyagin [3].

In this article are given the formulas for calculating the downwash upon the common/general/total formulation of the problem for the delta wing, which has supersonic leading edges.

Let the fine/thin slightly-curved delta wirg move in ideal

compressible liquid with low angle of attack and with certain angle of slip  $\beta_0$ . Let us consider that the basic motion of wing is rectilinear forward/progressive with the constant supersonic velocity U. Let us assume also that, besides basic motion, the wing completes small additional oscillation/vibrations.

Downwash is represented through the coefficients of the rotary derivatives [2, 3] and we will use the formulas of the calculation of these coefficients [2], which for the case of small Stroubal numbers take the form:

$$\theta(f)(x, y) = \frac{1}{\pi^{2}} \left\{ -\frac{1}{2} V. p. \int_{\chi(x)}^{z} \frac{f_{i}^{(t)}[\bar{\chi}(\eta), \eta] d\eta}{\sqrt{y - \eta^{2}} \sqrt{x - \bar{\chi}(\eta)}} + \int_{\chi(x)}^{z} \int_{\bar{\chi}(\eta)}^{z} \frac{1}{\sqrt{(x - \xi)(y - \eta)}} \frac{\partial^{2}}{\partial \xi \partial \eta} f_{i}^{(t)}(\xi, \eta) d\xi d\eta \right\}$$

$$(i = 1, 2; v = 1, 3, 4)$$

Formula (1) is written in the dimensionless characteristic coordinates xy, which are connected with the Cartesian coordinates x<sub>4</sub> 1, (Fig. 1) as follows:

$$z = \frac{2}{lk} z_1 - \frac{2}{l} y_1, \quad y = \frac{1}{lk} z_1 + \frac{2}{l} y_1. \tag{2}$$

where  $k=\sqrt{\hbar^2-1}$ ,  $\hbar=0/a$  - Each number, a - the speed of sound in the undisturbed flow, l - characteristic linear dimension (spread/scope) of wing.

Fage 46.

In formula (1) sign  $V.p. \int$  indicates the principal value of integral according to Hadamard [4], V = X(x) there is an equation of trailing wing edge, x = X(y) - the equation of the same wing edge, splved relatively by the variable x. Functions

 $f_i^{(i)}(x, y)$  (i = 1, 2, v = 1, 3, 4) are expressed by the formulas

$$f_{\gamma}^{(1)}(x, y) = -\int_{z_0 + z_1} \frac{B_{\gamma}^{(1)}(\xi, \eta) d\xi d\eta}{V(x - \xi)(y - \eta)} + A_{\gamma}^{(1)}(x, y)$$

$$f_{\gamma}^{(2)}(x, y) = \frac{\lambda}{8} \left( k + \frac{1}{k} \right) \int_{z_0 + z_1} B_{\gamma}^{(1)}(\xi, \eta) \frac{x - \xi + y - \eta}{V(x - \xi)(y - \eta)} d\xi d\eta + .$$

$$+ A_{\gamma}^{(2)}(x, y), \quad (v = 1, 3, 4)$$


$$(4)$$

where  $B_i^{(1)}(x, y)$  they are assigned by the condition of steady flow around of the wing and have following values [:2]:


$$B_1^{(1)}(x, y) = -1; \quad B_3^{(1)}(x, y) = -\frac{\lambda}{8}(x-y); \quad B_4^{(1)}(x, y) = -\frac{\lambda k}{8}(x+y)$$

λ= /2/S - aspect ratio, S - wing area.

Page 47.



Pig. 1.



Pig- 2-

The equations of wing edges will be (Fig. 1):

 $y=-i^2x$  - by rightist front/leading of edge;

 $y = -a_1^2x - by$  left front/leading of edge.

 $y = -\beta^{e}x + e = X(x) - a$  trailing edge,

where angular coefficients and number e are expressed as the geometric parameters of wing and the Mach number

$$e_{0}^{2} = -\frac{1+k \cdot g(\gamma - \beta_{0})}{1-k \cdot g(\gamma - \beta_{0})}, \quad e_{1}^{2} = -\frac{1-k \cdot tg(\gamma + \beta_{0})}{1+k \cdot tg(\gamma + \beta_{0})},$$

$$P = -\frac{1+k \cdot tg\beta_{0}}{1-k \cdot tg\beta_{0}}, \quad e = -\frac{8 \sqrt{1+ctg^{2}\beta_{0}}}{\lambda(1-k \cdot tg\beta_{0})}$$

By the interference waves, which proceed from point 0, wing is divided into three regions (I, II and I'), but tradling edge - to three cuts BC, CC' and C'A with different analytical expressions for potentials.

In formulas (3) and (4) functions  $A_i^{(i)}(x, y)$  are the values of the potentials of the disturbed velocities on the cuts indicated. For calculating these functions with small Stroubal numbers, we use the formulas, available in work [2].

Page 48.

For a delta wing when point M is arrange/located in region  $\sigma_2$  (Fig. 2), functions  $f_{ij}^{(n)}(x,y)$  have the following expressions:

$$f_{i}^{(1)}(x, y) = -\frac{\pi}{\beta} Z_{i}(x, y) + \sum_{j=0}^{1} g_{1, 1}^{(1, 1)} \left\{ Z_{j} \left[ \frac{\pi}{2} + (-1)^{j} \phi_{j} \right] - \frac{Q_{ij}}{\nabla} \left[ \frac{\pi}{2} + (-1)^{j} \phi_{j} \right] \right\}$$

$$= -\frac{Q_{ij}}{\nabla} \left[ \frac{\pi}{2} + (-1)^{j} \phi_{j} \right]$$

$$= -\frac{\lambda \pi}{8} \left[ \frac{e - \nabla x}{\beta} Z_{2}(x, y) - \frac{1 + 3\beta^{2}}{4\beta^{3}} Z_{2}^{2}(x, y) \right] + \frac{1}{2} \sum_{j=0}^{1} \sum_{n=0}^{2} g_{3, j}^{(1, j-n)} \left\{ Z_{j}^{2-n} y^{n} \left[ \frac{\pi}{2} + (-1)^{j} \phi_{j} \right] - \frac{1}{\nabla^{2}} Q_{1}^{n}(0) Q_{1, j}^{2-n} \left[ \frac{\pi}{2} + (-1)^{j} \phi_{j} \right] \right\} + \frac{1}{2} \sum_{n=0}^{1} \left[ b_{n, 1-n}^{(1)} x^{n} + \frac{1}{2} y^{\frac{3}{2}-n} + p_{n}^{(1)} (y - x)^{n} R(x, y) \right]$$

$$= -\frac{\lambda k \pi}{8} \left[ \frac{3\beta^{2} - 1}{4\beta^{3}} Z_{2}^{2}(x, y) + \frac{e + (1 - \beta^{2}) x}{\beta} Z_{2}(x, y) \right] + \frac{1}{2} \sum_{j=0}^{1} g_{3, j}^{(1, j-n)} \left\{ Z_{j}^{2-n} y^{n} \left[ \frac{\pi}{2} + (-1)^{j} \phi_{j} \right] - - \frac{1}{\nabla^{3}} Q_{1, j}^{2-n} Q_{1, j}^{n}(0) \left[ \frac{\pi}{2} + (-1)^{j} \phi_{j} \right] \right\} + \frac{1}{2} \left[ d_{n, 1-n}^{(1)} x^{n} x^{n} + \frac{1}{2} y^{\frac{3}{2}-n} + q_{1, n}^{(1)} (y - x)^{n} R(x, y) \right]$$

$$= f_{1}^{(2)} (x, y) = -\frac{\lambda M^{2} (1 - \beta^{2}) \pi}{2} Z_{1}^{2} (x, y) + \frac{\lambda M^{2} (1 - \beta^{2}) \pi}{2} Z_{2}^{2} (x, y) + \frac{\lambda M^{2} ($$

$$+ \sum_{j=0}^{1} g_{1,j}^{(2,2)} \left\{ Z_{j}^{2} \left[ \frac{\pi}{2} + (-1)^{j} \psi_{j} \right] - \frac{Q_{1,j}^{2}}{\nabla^{2}} \left[ \frac{\pi}{2} + (-1)^{j} \overline{\psi}_{j} \right] \right\} + \frac{1}{n=0} \left[ a_{n,1-n}^{(2)} x^{n+\frac{1}{2}} y^{\frac{3}{2}-n} + r_{n}^{(2)} (y-x)^{n} R(x,y) \right]$$

$$+ \sum_{n=0}^{1} \left[ a_{n,1-n}^{(2)} x^{n+\frac{1}{2}} y^{\frac{3}{2}-n} + r_{n}^{(2)} (y-x)^{n} R(x,y) \right]$$

$$+ \sum_{n=0}^{1} \sum_{n=0}^{1} g_{3,j}^{(2,3-n)} \left\{ Z_{j}^{3-n} y^{n} \left[ \frac{\pi}{2} + (-1)^{j} \psi_{j} \right] - \frac{Q_{1}^{n}(0) Q_{1}^{3-n}}{\nabla^{3}} \left[ \frac{\pi}{2} + (-1)^{j} \overline{\psi}_{j} \right] \right\} +$$

$$+ \sum_{n=0}^{2} \left[ b_{n,2-n}^{(2)} x^{n+\frac{1}{2}} y^{\frac{5}{2}-n} + p_{n}^{(2)} (y-x)^{n} R(x,y) \right]$$

$$+ \sum_{n=0}^{2} \left[ b_{n,2-n}^{(2)} x^{n+\frac{1}{2}} y^{\frac{5}{2}-n} + p_{n}^{(2)} (y-x)^{n} R(x,y) \right]$$

$$+ \sum_{n=0}^{1} \sum_{n=0}^{1} g_{4,j}^{(2,3-n)} \left\{ Z_{j}^{3-n} y^{n} \left[ \frac{\pi}{2} + (-1)^{j} \psi_{j} \right] - \frac{Q_{1,j}^{3-n} Q_{1}^{n}(0)}{\nabla^{3}} \left[ \frac{\pi}{2} + (-1)^{j} \overline{\psi}_{j} \right] \right\} +$$

$$+ \sum_{n=0}^{2} \left[ a_{n,2-n}^{(3)-n} Q_{1}^{n}(0) \left[ \frac{\pi}{2} + (-1)^{j} \overline{\psi}_{j} \right] \right\} +$$

$$+ \sum_{n=0}^{2} \left[ a_{n,2-n}^{(3)-n} x^{n+\frac{1}{2}} y^{\frac{5}{2}-n} + q_{n}^{(3)} (y-x)^{n} R(x,y) \right]$$

$$(10)$$

In formulas (5)-(10) and for further calculations, for purpose of a reduction in the recording, are introduced the following designations of the functions

$$Z_{j} = y + \alpha_{j}^{2}x, \quad Z_{1}(x, y) = y + \beta^{2}x - e, \quad Z_{2} = y + \beta^{3}x + e$$

$$Q_{0, j} = Q_{0}(y, \alpha_{j}) = \Delta_{j}y + \alpha_{j}^{2}e, \quad \overline{\chi}(y) = \frac{1}{\beta^{2}}(e - y)$$

$$Q_{1, j} = Q_{1}(x, y, \alpha_{j}) = \Delta_{j}(y - x) + (1 + \alpha_{j}^{2})e$$

$$Q_{1}(0) = Q_{1}(x, y, 0) = \beta^{2}(y - x) + e$$

$$Q_{2, j} = Q_{2}(x, \alpha_{j}) = -\Delta_{j}x + e$$

$$Q_{3, j} = Q_{3}(x, \alpha_{j}) = -\Delta_{j}x + (\alpha_{j}^{2} + \beta^{2})e$$

$$Q_{j} = Q(x, \alpha_{j}) = \alpha_{j}^{2}(37\beta^{2}x + 14e)$$

$$\psi_{1, j} = \psi_{1}(x, y, \alpha_{j}) = \arcsin \frac{y - \alpha_{j}^{2}x}{y + \alpha_{j}^{2}x}, \quad \psi_{0, j} = \overline{\psi}_{0, j} = \frac{\pi}{2}$$

$$\overline{\psi}_{1, j} = \overline{\psi}_{2}(x, y, \alpha_{j}) = \arcsin \frac{(\beta^{2} + \alpha_{j}^{2})(y - x) + (1 - \alpha_{j}^{2})e}{\Delta_{j}(y - x) + (1 + \alpha_{j}^{2})e}$$

$$C(y, \alpha_{j}) = \arcsin \frac{(\alpha_{j}^{2} + \beta^{2})y - \alpha_{j}^{2}e}{\Delta_{j}y + \alpha_{j}^{2}e}$$

$$R(x, y) = \sqrt{e^{2} + (\beta^{2} - 1)e(y - x) - \beta^{2}(y - x)^{2}}$$

$$R_{1}(x, y, \alpha_{j}) = \operatorname{arctg} \sqrt{\frac{y + \beta^{2}x - e}{(e - y)(y + \alpha_{j}^{2}x)}}$$

$$R_{2}(x, y) = \operatorname{arctg} \sqrt{\frac{y + \beta^{2}x - e}{e - y}}$$

Fage 49.

Are given below the values of coefficients, which participate in the calculations which depend only on the geometric wing characteristics ( $\alpha_0$ ,  $\alpha_1$ ,  $\beta$ ,  $\lambda$ ) and of Mach runber.

$$\begin{aligned}
 & \eta_1 = a_0^2 a_1^2, \quad \gamma_2 = a_0^2 - a_1^2, \quad \gamma_3 = a_0^2 + a_1^2, \quad \Delta_j = \beta^2 - a_j^2, \quad \nabla = 1 + \beta^2 \\
 & g_{1,j}^{(1,j)} = \frac{1}{a_j}, \quad g_{1,j}^{(2,2)} = -\frac{\lambda M^2}{32k} \frac{1 + a_j^2}{a_j^3}, \quad g_{3,j}^{(1,j)} = -\frac{\lambda (1 + a_j^2)}{8a_j^3} \\
 & g_{3,j}^{(1,1)} = \frac{\lambda (3 + a_j^2)}{32 a_j^3}, \quad g_{3,j}^{(2,2)} = \frac{\lambda^2 M^2}{256k} \frac{(1 + a_j^2)^2}{a_j^5}, \quad g_{3,j}^{(2,3)} = -\frac{\lambda^2 M^2}{512k} \frac{(1 + a_j^2)^2}{a_j^5} \\
 & g_{4,j}^{(1,1)} = -\frac{\lambda k (1 - a_j^2)}{8a_j^3}, \quad g_{4,j}^{(1,2)} = \frac{\lambda k}{32} \frac{3 - a_j^2}{a_j^3}, \quad g_{4,j}^{(2,2)} = \frac{\lambda^2 M^2}{256} \frac{a_j^4 - 1}{a_j^5} \\
 & g_{4,j}^{(1,1)} = -\frac{\lambda^2 M^2 (3a_j^4 - 4a_j^2 - 3)}{1536 a_j^5}, \quad (j = 0, 1) \\
 & g_{1,0}^{(2)} = \frac{\gamma_2 M^2 \lambda}{16k}, \quad a_{0,1}^{(2)} = \frac{\lambda \gamma_2 M^2}{16 \gamma_1 k}, \quad b_{1,0}^{(1)} = \frac{\lambda \gamma_2}{16}, \quad b_{0,1}^{(1)} = -\frac{\lambda \gamma_2}{16 \gamma_1} \\
 & b_{2,0}^{(2)} = \frac{\lambda^2 M^2}{256k} \gamma_2 (2 + \gamma_3), \quad b_{1,0}^{(2)} = -\frac{\lambda^2 M^2 \gamma_2}{384 k \gamma_1} (1 - \gamma_1) \\
 & b_{0,2}^{(2)} = -\frac{\lambda^2 M^2 \gamma_2}{256k \gamma_1^2} (\gamma_3 + 2\gamma_1), \quad d_{1,0}^{(1)} = \frac{\lambda k \gamma_2}{16}, \quad d_{0,1}^{(1)} = \frac{\lambda k \gamma_2}{16 \gamma_1} \\
 & d_{0,2}^{(2)} = \frac{\lambda^2 M^2 \gamma_2}{768} \gamma_2 (4 - 3\gamma_3), \quad d_{1,1}^{(2)} = -\frac{\lambda^2 M^2 \gamma_2}{384 \gamma_1} (1 - \gamma_1) e \\
 & p_{1,0}^{(1)} = \frac{\lambda \gamma_2 (\beta^2 + \gamma_1)}{16 \gamma_2 \gamma_2} \\
 & h_{1,0}^{(2)} = \frac{\lambda^2 M^2 \gamma_2}{768 \gamma_1^2} (4\gamma_1 - 3\gamma_2), \quad p_{0,0}^{(1)} = \frac{\lambda \gamma_2}{16 \gamma_1^2} (1 - \gamma_1) e

 \end{aligned}$$

$$p_0^{(3)} = \frac{\lambda^3 \gamma_2 M^3}{768 \gamma_1^2 k V^3} \left[ -3 \gamma_1^3 (2 + \gamma_4) + 2 \gamma_1 (1 - \gamma_5) + 3 (2 \gamma_2 + \gamma_4) \right] e^3$$

$$p_1^{(2)} = -\frac{\lambda^2 M^2 \gamma_5}{768 k \gamma_1^2 V^3} \left[ 6 \gamma_1^2 (2 + \gamma_4) + 2 (\beta^2 - 1) (1 - \gamma_1) \gamma_1 + 6 \beta^2 (2 \gamma_1 + \gamma_4) \right] e$$

$$p_2^{(2)} = -\frac{\lambda^2 M^2 \gamma_5}{768 k \gamma_1^2 V^3} \left[ 3 \gamma_1^2 (2 + \gamma_4) + 2 \beta^2 \gamma_1 (1 - \gamma_1) - 3 (2 \gamma_1 + \gamma_4) \beta^4 \right]$$

$$q_0^{(1)} = \frac{\lambda k \gamma_5}{16 \gamma_1 V^2} (1 + \gamma_1) e, \quad q_1^{(1)} = \frac{\lambda k \gamma_5}{16 \gamma_1 V^2} (\beta^2 - \gamma_1)$$

$$q_0^{(2)} = -\frac{\lambda^2 M^2 \gamma_5}{768 \gamma_1^2 V^3} \left[ \gamma_1^2 (4 - 3 \gamma_3) - 2 \gamma_1 (1 + \gamma_1) + (4 \gamma_1 - 3 \gamma_3) \right] e^3$$

$$q_1^{(2)} = \frac{\lambda^2 M^2 \gamma_5}{768 \gamma_1^2 V^3} \left[ 2 \gamma_1^2 (4 - 3 \gamma_3) - 2 \gamma_1 (1 + \gamma_1) (1 - \beta^2) - 2 \beta^2 (4 \gamma_1 - 3 \gamma_3) \right] e$$

$$q_2^{(2)} = -\frac{\lambda^2 M^2 \gamma_5}{768 \gamma_1^2 V^3} \left[ \gamma_1^2 (4 - 3 \gamma_3) + 2 \beta^2 \gamma_1 (1 + \gamma_1) + \beta^4 (4 \gamma_1 - 3 \gamma_3) \right]$$

$$r_0^{(2)} = -\frac{\lambda M^2}{16k} \gamma_2 \frac{1 + \gamma_1}{V^2 \gamma_1} e, \quad r_1^{(2)} = -\frac{\lambda M^2}{16k} \gamma_5 \frac{\mu_5 - \gamma_1}{V^2 \gamma_1}$$

$$w_1 = 3e \beta^2 - 14 a_1^2$$

Page 50.

Substituting in formula (2) of the value of functions  $f_i^{(n)}(x, y)$  from (5)-(.10) and performing integration, we will obtain expressions for  $\theta_i^{(n)}(x, y)$  in the following form:

$$\theta_{1}^{(1)}(x, y) = -1 + \frac{1}{\pi^{2}} \sum_{j=0}^{1} g_{1,j}^{(1,j)} \left\{ F_{0,j}^{(0,j)}(x, y) + (-1)^{j} F_{1,j}^{(0,j)}(x, y) - \frac{1}{\nabla} \left[ L_{0,j}^{(0,j)}(x, y) + (-1)^{j} L_{1,j}^{(0,j)}(x, y) \right] \right\}$$

$$\theta_{3}^{(1)}(x, y) = \frac{\lambda}{8} \left\{ \frac{1 + 3\beta^{2}}{2\beta^{2}} Z_{2}(x, y) + (e - \nabla x) + \frac{1}{\pi^{2}} \sum_{j=0}^{1} \sum_{n=0}^{1} g_{3,j}^{(1,2-n)} \left\{ F_{0,j}^{(n,2-n)}(x, y) + (-1)^{j} F_{1,j}^{(n,2-n)}(x, y) - \frac{1}{\nabla^{2}} \left[ L_{0,j}^{(n,2-n)}(x, y) + (-1)^{j} L_{0,j}^{(n,2-n)}(x, y) \right] \right\} + \frac{1}{\pi^{2}} \sum_{n=0}^{1} \left\{ b_{n,1-n}^{(1)} H_{n,1-n}(x, y) + p_{n}^{(1)} M_{n}(x, y) \right\} + \frac{1}{\pi^{2}} \sum_{n=0}^{1} \left\{ b_{n,1-n}^{(1)} H_{n,1-n}(x, y) + p_{n}^{(1)} M_{n}(x, y) \right\} \right\}$$

$$\theta_{4}^{(1)}(x, y) = -\frac{\lambda}{8} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \sum_{n=0}^{\infty} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^{2} - 1}{4\beta^{2}} Z_{2}(x, y) + (1 - \beta^{2}) x + e \right\} + \frac{1}{\pi^{2}} \left\{ \frac{3\beta^$$

Fage 51.

$$+\frac{1}{\pi^2}\sum_{j=0}^{3}\sum_{n=0}^{1}g_{n,j}^{(1,2-n)}\left\{F_{0,j}^{(n,2-n)}(x,y)+(-1)^jF_{1,j}^{(n,2-n)}(x,y)\right] + \\ -\frac{1}{\tau^2}\left[L_{0,j}^{(n,2-n)}(x,y)+(-1)^jL_{1,j}^{(n,n-2)}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{3}\left[d_{n,1-n}^{(1)}H_{n,1-n}(x,y)+q_{n}^{(1)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{j=0}^{1}g_{1,j}^{(2)}\left\{F_{0,j}^{(0,2)}(x,y)+(-1)^jF_{1,j}^{(0,2)}(x,y)+ \\ -\frac{1}{\tau^2}\sum_{j=0}^{1}\left[L_{0,j}^{(2)}(x,y)+(-1)^jL_{1,j}^{(0,2)}(x,y)\right]\right\} + \\ +\frac{1}{\pi^2}\sum_{n=0}^{1}\left[a_{n,1-n}^{(2)}H_{n,1-n}(x,y)+r_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{1}\left[a_{n,1-n}^{(2)}H_{n,1-n}(x,y)+r_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{1}\sum_{n=0}^{1}g_{3,j}^{(2,3-n)}\left\{F_{0,j}^{(n,j-n)}(x,y)+(-1)^jF_{1,j}^{(n,j-n)}(x,y)- \\ -\frac{1}{\tau^2}\left[L_{0,j}^{(n,3-n)}(x,y)+(-1)^jL_{1,j}^{(n,3-n)}(x,y)\right]\right\} + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[b_{n,2-n}^{(2)}H_{n,2-n}(x,y)+p_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[b_{n,2-n}^{(2)}H_{n,2-n}(x,y)+p_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[L_{0,j}^{(n,3-n)}(x,y)+(-1)^jL_{1,j}^{(n,3-n)}(x,y)+ \\ -\frac{1}{\eta^3}\left[L_{0,j}^{(n,3-n)}(x,y)+(-1)^jL_{1,j}^{(n,3-n)}(x,y)\right]\right\} + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[L_{0,j}^{(n,3-n)}(x,y)+(-1)^jL_{1,j}^{(n,3-n)}(x,y)\right]\right\} + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right]\right\} + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right]\right\} + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right]\right\} + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}(x,y)\right] + \\ +\frac{1}{\pi^2}\sum_{n=0}^{2}\left[d_{n,2-n}^{(2)}H_{n,2-n}(x,y)+q_{n}^{(2)}N_{n}$$

PAGE 12

Page 52.

In formulas (11) - (16) for downwashes  $\theta_{ij}^{(j)}(x, y)$  enter the functions

$$F_{l,j}^{(n,m)}(x, y) = \frac{\partial}{\partial y} \int_{\chi(x)}^{x} \frac{d\eta}{\sqrt{y-\eta}} \frac{\partial}{\partial x} \int_{\overline{\chi}(\eta)}^{x} \frac{\eta^{n} Z_{l}^{n}}{\sqrt{x-\xi}} \phi_{l}(\xi, \eta, a_{l}) d\xi$$

$$(n = 0, 1; m = 1, 2, 3; l = 0, 1)$$

$$L_{l,j}^{(n,m)}(x, y) = \frac{\partial}{\partial y} \int_{\chi(x)}^{y} \frac{d\eta}{\sqrt{y-\eta}} \frac{\partial}{\partial x} \times$$

$$\times \int_{\overline{\chi}(\eta)}^{x} \frac{Q_{1}^{n}(\xi, \eta, 0) Q_{1}^{m}(\xi, \eta, a_{j})}{\sqrt{x-\xi}} \phi_{l}(\xi, \eta, a_{j}) d\xi$$

$$H_{n,m}(x, y) = \frac{\partial}{\partial y} \int_{\chi(x)}^{y} \frac{\eta^{m} \sqrt{\eta} d\eta}{\sqrt{y-\eta}} \frac{\partial}{\partial x} \int_{\overline{\chi}(\eta)}^{x} \frac{\xi^{n} \sqrt{\xi} d\xi}{\sqrt{x-\xi}}, (n, m = 0, 1, 2)$$

$$N_{n}(x, y) = \frac{\partial}{\partial y} \int_{\chi(x)}^{y} \frac{d\eta}{\sqrt{y-\eta}} \frac{\partial}{\partial x} \int_{\overline{\chi}(\eta)}^{x} \frac{(\eta - \xi)^{n} R(\xi, \eta) d\xi}{\sqrt{x-\xi}}$$

which, after the execution of actions in right sides, take the form

$$F_{0,j}^{(0,j)}(x, y, \alpha_{j}) = \frac{\pi^{2} (\alpha_{j}^{2} + \beta^{2})}{4 \beta \alpha_{j}}$$

$$F_{0,j}^{(0,j)}(x, y, \alpha_{j}) = \frac{\pi^{2}}{8\beta^{3}} \left\{ \left[ 4 (2\beta^{2} + \Delta_{j}) \alpha_{j}^{2} - 3\Delta_{j}^{2} \right] y + \beta^{2} \left[ \Delta_{j}^{2} + 4\alpha_{j}^{2} (\beta^{2} + \alpha_{j}^{2}) \right] x - \Delta_{j}^{2} e \right\}$$

$$F_{0,j}^{(0,j)}(x, y, \alpha_{j}) = \frac{3\pi^{2}}{32\beta^{5}} \left\{ -\Delta_{j}^{3} (4y^{2} + Z_{3}^{2}) - 4\alpha_{j}^{2} \Delta_{j}^{2} e (2y + Z_{3}^{2}) + 8\alpha_{j}^{4} \Delta_{j} e^{2} + 6\alpha_{j}^{2} Q_{0,j}^{2} + 4\alpha_{j}^{2} \beta^{2} Q_{0,j} Q_{2,j} + (19) + 6\alpha_{j}^{2} \beta^{4} Q_{2,j}^{2} + 24\alpha_{j}^{2} \beta^{2} Z_{2}(x, y) (\beta^{2} Q_{2,j} + 3Q_{0,j}) + 16\alpha_{j}^{4} \beta^{2} Z_{2}^{2}(x, y) \right\}$$

$$F_{0,j}^{(1,1)}(x, y, \alpha_{j}) = -\frac{\pi^{2}}{8\beta} \left( 3\Delta_{j} y - \beta^{2} \Delta_{j} x + \beta^{2} e - 3\alpha_{j}^{2} e \right) + \frac{\pi^{2} \alpha_{j}^{2}}{4\beta} \left[ 2y + Z_{1}(x, y) \right]$$

$$F_{0,j}^{(1,2)}(x, y, \alpha_{j}) \approx -\frac{3\pi^{2} \Delta_{j}}{32\beta^{3}} \left( 4y^{2} + Z_{3}^{2} \right) + \frac{\pi^{2} \alpha_{j}^{2}}{4\beta^{3}} Z_{2}(x, y) \times \left[ (\alpha_{j}^{2} - 4\Delta_{j}) \beta^{2} x + (5\beta^{2} + \Delta_{j}) y + (\beta^{2} + 2\alpha_{j}^{2}) e \right] +$$

Page 53.

$$\begin{aligned} &+\frac{\pi^2 a_1^2}{4 \beta^2} \left[ 4 \beta^4 (\Delta_1 x - e) x + 5 a_2^2 e^2 \right] \\ &L_{0,1}^{(0,1)}(x, y, a_1) = -\frac{\Delta_1 \pi^2}{4 a_1 \beta} + \frac{\pi^2 \Delta_1}{2 \sqrt{a_1 \beta}} (y + \beta^2 x - e) \end{aligned}$$
(22)
$$L_{0,1}^{(0,1)}(x, y, a_1) = -\frac{\pi^2 \Delta_1}{8 \beta^3} \left\{ \Delta_1 (3 \nabla^2 + 12 \nabla - 8) y - \Delta_1 (\nabla^2 - 4 \nabla + 8) \beta^2 x + \left[ \nabla^2 (\beta^2 - 5 a_1^2) + 4 \beta^2 (1 - \beta^2) + 4 a_1^2 (1 + 3 \beta^2) \right] e \right\}$$
(23)
$$L_{0,1}^{(0,3)}(x, y) = -\frac{3\pi^2 \nabla^2 \Delta_1}{32 \beta^3} \left\{ \Delta_2^2 (4 y^2 + Z_3^2) + 4 a_1^2 \Delta_1 e (2 y + Z_3) - 8 a_1^4 e^2 \right\} - \frac{3\pi^2}{16 \beta^3} \Delta_1 \left[ 8 \beta^4 Q_0^2 (y, a_1) + 2 \beta^2 (3 + 2 \beta^2) \Delta_1 Q_0 (y, a_1) \times X \right] \\ &\times Z_1(x, y) + \Delta_1^2 (3 \beta^4 + 4 \beta^2 + 3) Z_2^2(x, y) + 8 \beta^4 Q_0 (y, a_1) Q_1(x, a_1) + 8 \beta^4 \nabla Q_2^2(x, a_1) + 2 \beta^2 (2 \nabla - 1) (\nabla + 3) \Delta_1 Q_2(x, a_1) Z_1(x, y) \right\}$$
(24)
$$L_{0,1}^{(1,1)}(x, y) = -\frac{\pi^2 \nabla^2}{8 \beta} \left[ \Delta_1 (2 y + Z_1) - 2 a_2^3 e \left[ -\frac{\pi^2}{4 \beta} Z_1(x, y) \times X \right] X \right] \\ &\times \left[ (3 \nabla - 2) \Delta_1 y - \Delta_1 \nabla \beta^2 x + (\beta^4 + a_2^2 \beta^2 + \Delta_1) e \right]$$
(25)
$$L_{0,1}^{(1,2)}(x, y) = -\frac{\pi^2 \nabla^2}{32 \beta^3} \left[ 3 \Delta_1^2 \left[ 4 y^2 + Z_2^2 \right] + 8 \Delta_1 a_2^2 e \left[ 2 y + Z_2 \right] - 8 a_1^4 e^3 \right] - \frac{3\pi^2}{16 \beta^3} \Delta_2^3 \nabla (4 - \nabla) Z_2^2(x, y) - \frac{\pi^2}{2 \beta^3} \left[ \Delta_1 (2 \nabla - 1) \times X \right] \\ &\times \left[ 3 \beta^9 Q_{1,1} - a_1^2 \nabla e \right] Z_1(x, y) + 3 \beta^4 \left[ Q_0(y, a_1) - \Delta_1 Z_2(x, y) \right] Q_0(y, a_1) + 3 \beta^4 \left[ Q_{1,1} + \beta^9 Q_0(x, a_1) \right] Q_1(x, a_1) - a_2^2 e \nabla \left[ Q_{1,1} + \nabla Q_0(x, a_1) \right] (26)$$

$$F_{1,1}^{(0,1)}(x, y, a_1) = -\frac{1}{2 \beta} V.p. \int_{1/2}^{p} \frac{Q_0(\eta_1, a_1) C(\eta_1, a_1) d\eta}{\sqrt{y - \eta} V Z_1(x, \eta)} - \frac{2}{1/2 \beta} V + \beta^2 x - e} - f_1 + \frac{a_2^2}{\beta} \int_{1/2}^{p} \frac{R_1(x, \eta_1, a_1) d\eta}{\sqrt{y - \eta} V \gamma + a_2^2 x} + a_1 \int_{1/2}^{p} \frac{R_2(x, \eta) d\eta}{\sqrt{\eta} (y - \eta)} - \frac{2}{1/2 \beta} V + \beta^2 x - e} - \frac{1}{2 \beta} \left[ (e_1^2 + 2 \beta^2) J_0 - a_2^2 \left[ 3 \chi(x) + e \right] J_1 + 3 a_2^2 e \chi(x) J_0^2 + \frac{2}{\beta} \chi(x)$$

Rage 54.

$$+\frac{2a_{j}^{2}}{\beta^{9}} \int_{I(z)}^{z} [Q_{0}(\eta, a_{j}) + \mathbf{Z}_{0}(z, \eta)] \frac{C(\eta, a_{j})}{\sqrt{y-\eta}} \frac{A}{\sqrt{Z_{0}(z, \eta)}} + Q_{0}^{2} \frac{10\eta + 3a_{j}^{2}x}{\sqrt{(y-\eta)\eta}} R_{0}(x, \eta) d\eta - \\ -8a_{j} \int_{I(z)}^{z} \sqrt{\frac{\eta + a_{j}^{2}x}{y-\eta}} R_{1}(x, \eta, a_{j}) d\eta$$

$$F_{1,j}^{(a_{j},5)}(x, y, a_{j}) = -\frac{1}{2\beta^{9}} V.p. \int_{I(z)}^{z} \frac{Q_{0}^{3}(\eta, a_{j}) C(\eta, a_{j}) d\eta}{Vy - \eta^{3}VZ_{0}(x, \eta)} + \\ +\frac{a_{j}^{2}}{\beta^{3}} \int_{I(z)}^{z} [3Q_{0}^{2}(\eta, a_{j}) + 12\beta^{9}Q_{0}(\eta, a_{j}) Z_{1}(x, \eta) + \\ +8a_{j}^{2}\beta^{2}Z_{2}^{2}(x, \eta)] \frac{C(\eta, a_{j}) d\eta}{Vy - \eta VZ_{1}(x, \eta)} + \\ +\frac{a_{j}}{80} \int_{I(z)}^{z} [880\eta^{2} + 960a_{j}^{2}x\eta + 165a_{j}^{4}x^{2}] \frac{R_{1}(x, \eta)}{V\eta(y-\eta)} d\eta - \\ -16a_{j} \int_{I(z)}^{z} \sqrt{\frac{(\eta + a_{j}^{2}x)^{3}}{y - \eta}} R_{1}(x, \eta) d\eta + \frac{2ea_{j}^{2}}{\beta^{3}} [(3\beta^{2} + a_{j}^{2}) J_{z} + \\ + [3a_{j}^{2}e - (3\beta^{2} + 5a_{j}^{2})X(x)] J_{1} - a_{j}^{2}X(x) (4\beta^{2}x - e) J_{0}] + \\ +\frac{a_{j}}{30\beta^{4}} [-5\omega_{j}J_{3} + [4\omega_{j}(2e - \beta^{3}x) - 3\Omega_{j}(x)] J_{2} - [3\omega_{j}eX(x) - \\ -2\Omega_{j}(x)(2e - \beta^{3}x)] J_{1} - e\Omega_{j}(x)X(x)J_{0}] + \\ +\frac{a_{j}}{80} [176J_{3} + 320a_{j}^{2}xJ_{2} + 165a_{j}^{4}x^{2}J_{1}] - \frac{16}{5}a_{j}[J_{3} + \\ + 2a_{j}^{2}xJ_{2} + a_{j}^{4}x^{2}J_{3}] - \frac{16a_{j}^{3}e}{5\beta^{4}} ((a_{j}^{2} + \beta^{3})J_{2} + [a_{j}^{2}\beta^{3}x - e)X(x)J_{0}]$$

$$F_{1,j}^{(1,1)}(x, y, a_{j}) = -\frac{1}{2\beta} V.p. \int_{I(z)}^{z} \frac{\eta Q_{0}(\eta, a_{j}) C(\eta, a_{j})}{Vy - \eta^{-3}VZ_{1}(x, \eta)} d\eta + \\ +3a_{j} \int_{I(z)}^{(1,1)} \frac{\eta Q_{0}(\eta, a_{j}) C(\eta, a_{j})}{y - \eta^{-3}VZ_{1}(x, \eta)} d\eta + \\ +3a_{j} \int_{I(z)}^{(1,1)} \frac{\eta Q_{0}(\eta, a_{j}) C(\eta, a_{j})}{y - \eta^{-3}VZ_{1}(x, \eta)} d\eta + \\ +3a_{j} \int_{I(z)}^{(1,1)} \frac{\eta Q_{0}(\eta, a_{j}) C(\eta, a_{j})}{y - \eta^{-3}VZ_{1}(x, \eta)} d\eta + \\ +3a_{j} \int_{I(z)}^{(1,1)} \frac{\eta Q_{0}(\eta, a_{j}) C(\eta, a_{j})}{y - \eta^{-3}VZ_{1}(x, \eta)} d\eta + \\ +3a_{j} \int_{I(z)}^{(1,1)} \frac{\eta Q_{0}(\eta, a_{j}) C(\eta, a_{j})}{y - \eta^{-3}VZ_{1}(x, \eta)} d\eta + \\ +3a_{j} \int_{I(z)}^{(1,1)} \frac{\eta Q_{0}(\eta, a_{j}) C(\eta, a_{j})}{\eta Q_{0}(\eta, a_{j})} d\eta + \\ \frac{\eta Q_{0}(\eta, a_{j})}{\eta Q_{0}(\eta, a_{j})} \frac{\eta Q_{0}(\eta, a_{j})}{\eta Q_{0}(\eta, a_{j})} d\eta + \\ \frac{\eta Q_{0}(\eta, a_{j})}{\eta Q_{0}(\eta, a_{j})} \frac{\eta Q_{$$

Fage 55.

$$\begin{aligned}
&+\frac{a_{1}^{2}}{2}\int_{\chi(x)}^{\infty}\frac{3\eta+2\theta^{2}x-2a}{Vy-\eta}\frac{C(x,a_{1})d\eta-a_{1}f_{2}-}{V(y-\eta)(\eta+a_{2}^{2}x)}d\eta \\
&-2a_{1}\int_{\chi(x)}^{\pi}\frac{(3\eta+2a_{1}^{2}x)R_{1}(x,\eta_{1}|x_{1})}{V(y-\eta)(\eta+a_{2}^{2}x)}d\eta \\
&F_{1,j}^{(1,j)}(x,y,a_{1}) = -\frac{1}{2\beta^{2}}V.p.\int_{\chi(x)}^{\pi}\frac{\eta Q_{0}^{2}(\eta_{1}|x_{1})}{\sqrt{y-\eta^{2}}VZ_{1}(x,\eta)}d\eta + \\
&+\frac{2a_{1}^{2}}{\beta^{2}}\int_{\chi(x)}^{\pi}\left[Q_{0}(\eta_{1}|a_{1})+2\beta^{2}Z_{1}(x,\eta)\right]\frac{\eta C(\eta_{1}|a_{1})d\eta}{V(y-\eta)Z_{1}(x,\eta)} + \\
&+\frac{a_{2}}{6}\int_{\chi(x)}^{\pi}\left[5\eta+2a_{2}^{2}x\right]V\frac{\eta+a_{2}^{2}x}{y-\eta}R_{1}(x,\eta_{1}|a_{1})d\eta + \\
&+\frac{4a_{2}^{2}}{\beta^{2}}\int_{\chi(x)}^{\pi}\left[Q_{0}(\eta_{1}|a_{1})Z_{2}(x,\eta)+\frac{2}{3}a_{2}^{2}Z_{2}^{2}\right]V\frac{Z_{1}(x,\eta)}{y-\eta}C(\eta_{1}|a_{1})d\eta + \\
&+\frac{4a_{2}^{2}}{\beta^{2}}\int_{\chi(x)}^{\pi}\left[Q_{0}(\eta_{1}|a_{1})Z_{2}(x,\eta)+\frac{2}{3}a_{2}^{2}Z_{2}^{2}\right]V\frac{Z_{1}(x,\eta)}{y-\eta}C(\eta_{1}|a_{1})d\eta - \\
&-\frac{a_{1}}{\beta^{2}}\left\{(a_{1}^{2}+\beta^{2})J_{2}-a_{1}^{2}[a+\chi(x)]J_{2}+2a_{2}^{2}e\chi(x)J_{1}\right\} \qquad (31)
\end{aligned}$$

$$L_{1,j}^{(0,j)}(x,y,a_{j}) = -\frac{\nabla}{2\beta}V.p.\int_{\chi(x)}^{\pi}\frac{Q_{0}(\gamma_{1}|a_{1})C(\eta_{1}|a_{1})d\eta}{V(y-\eta)^{2}Z_{1}(x,\eta)} - \\
&-\frac{\Delta_{1}}{\beta}\int_{\chi(x)}^{\pi}\frac{C(\eta_{1}|a_{1})d\eta}{V(y-\eta)}\frac{d\eta}{Z_{2}(x,\eta)} - \frac{3\nabla-2}{\gamma}a_{1}ef_{0}+ \\
&+\frac{2\beta^{2}}{\Delta}Q_{2}(x,a_{1})J_{-1}-\frac{\Delta_{1}}{a_{1}\beta}\int_{\chi(x)}^{\pi}\frac{G_{1}^{(1)}(x,\eta_{1}|a_{1})d\eta}{V(y-\eta)}d\eta + \\
&+\frac{a_{1}ey}{2\beta}\int_{\chi(x)}^{\pi}\frac{G_{1}^{(0)}(x,\eta_{1}|a_{1})d\eta}{V(y-\eta)}d\eta - \frac{32}{\gamma}Z_{1}(x,\eta)d\eta - \\
&+\frac{a_{1}ey}{\beta}\int_{\chi(x)}^{\pi}\frac{G_{1}^{(0)}(x,\eta_{1}|a_{1})d\eta}{V(y-\eta)}d\eta - \frac{32}{\gamma}Z_{1}(x,\eta)d\eta - \\
&+\frac{a_{1}ey}{\beta}\int_{\chi(x)}^{\pi}\frac{G_{1}^{(0)}(x,\eta_{1}|a_{1})d\eta}{U(y-\eta$$

Page 56.

$$-\frac{2\Delta_{j}}{\beta^{3}}\int_{z(s)}^{z} [\beta^{a}Q_{a,j} + 2\eta\Delta_{j}Z_{a}(x,\eta)] \frac{C(\eta,\alpha_{j})d\eta}{\sqrt{y-\eta}} \frac{-2(x,\eta)}{\sqrt{Z_{a}(x,\eta)}} - \frac{4\alpha_{j}e}{3\eta} \left[ (3\eta - 2)\Delta_{j}f_{1} + [(3\eta - 2)e - \eta Q_{a}(x,\alpha_{j})] f_{0} + \frac{2\beta^{4}}{\Delta_{j}}Q_{2}^{2}(x,\alpha_{j})f_{-1} \right] - \frac{\eta\alpha_{j}e}{\beta^{3}} \left[ \gamma_{j}f_{1} + \alpha_{j}^{2}ef_{0} \right] + \frac{4}{\beta} \nabla\alpha_{j}\Delta_{j}e \int_{\chi(s)}^{z} \left\{ f_{-1}^{(1)}(p_{s}) - \frac{Q_{1}(x,\eta,\alpha_{j})}{2\alpha_{j}^{2}\eta e} G_{1}^{(1)}(x,\eta,\alpha_{j}) - \frac{\Delta_{j}G_{1}^{(2)}(x,\eta,\alpha_{j})}{6\eta\alpha_{j}^{2}e} \right\} \frac{d\eta}{\sqrt{y-\eta}} - \frac{\alpha_{j}\eta e}{2\beta} \int_{\chi(s)}^{z} \left\{ 2\Delta_{j}f_{0} - Q_{1}(x,\eta,\alpha_{j}) \times X G_{2}^{(0)}(x,y,\alpha_{j}) - \Delta_{j}G_{2}^{(1)}(x,\eta,\alpha_{j}) \right\} \frac{d\eta}{\sqrt{y-\eta}} \left( 33 + \Delta_{1,j}^{2}(x,y,\alpha_{j}) \right) = -\frac{\nabla^{2}}{2\beta^{2}} V \cdot p \cdot \int_{\chi(s)}^{z} \frac{Q_{0}^{2}(\eta,\alpha_{j})}{\sqrt{y-\eta}^{3}VZ_{1}(x,\eta)} d\eta - \frac{\Delta_{j}}{3\beta^{3}} \int_{\chi(s)}^{z} \left[ 3\beta^{4}Q_{1}^{2}(x,\eta,\alpha_{j}) + 6\beta^{2}\Delta_{j}(2\gamma - 1) Q_{1}(x,\eta,\alpha_{j}) Z_{2}(x,\eta) + \Delta_{j}^{2}(4\gamma - 1) Z_{2}^{2}(x,\eta) \right] \frac{C(\eta,\alpha_{j})d\eta}{\sqrt{y-\eta}VZ_{2}(x,\eta)} - \frac{2\alpha_{j}e}{5\gamma\beta^{2}} \left\{ 15\beta^{4} + 10\beta^{2} + 3) [\Delta_{j}^{2}f_{2} + 2\alpha_{j}^{2}e\Delta_{j}f_{1} + \alpha_{j}^{4}e^{2}f_{0}] - \beta^{2}(15\beta^{4} - 10\beta^{2} - 1) Q_{2}(x,\alpha_{j})[\Delta_{j}f_{1} + \alpha_{j}^{2}eJ_{0}] + 4\beta^{4}(1 - 5\beta^{2})Q_{2}^{2}(x,\alpha_{j})f_{0} + \frac{8\beta^{6}}{\Delta_{j}}Q_{2}^{3}(x,\alpha_{j})f_{-1} \right\} - \frac{\alpha_{j}e}{\beta^{2}} \left\{ \Delta_{j}^{2}(3\gamma - 2) f_{2} + 2\Delta_{j} \left[ 2\beta^{2}Q_{1}(x,0,\alpha_{j}) - \nabla\Delta_{j}X(x) \right] f_{1} + (\beta^{2}Q_{1}(x,\eta,\alpha_{j}))[Q_{1}(x,\eta,\alpha_{j}) + 20\gamma_{2}^{2}\Delta_{j}A_{j}ef_{-1}^{2}(p_{3}) - 15Q_{1}^{2}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}) + 10\Delta_{j}Q_{1}(x,\eta,\alpha_{j}) G_{1}^{(2)}(x,\eta,\alpha_{j}) f_{0} + \frac{\Delta_{j}}{5\alpha_{j}}G_{1}^{(2)}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}) - \frac{\eta\alpha_{j}e}{2\beta} \int_{\xi}^{\xi} \left[ 4\Delta_{j}Q_{1}(x,\eta,\alpha_{j}) f_{0} + + 4\Delta_{j}^{2}f_{1} - Q_{1}^{2}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}) - \frac{\eta\alpha_{j}e}{2\beta} \int_{\xi}^{\xi} \left[ 4\Delta_{j}Q_{1}(x,\eta,\alpha_{j}) f_{0} + + 4\Delta_{j}^{2}f_{1} - Q_{1}^{2}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}) - \frac{\eta\alpha_{j}e}{2\beta} \int_{\xi}^{\xi} \left[ 4\Delta_{j}Q_{1}(x,\eta,\alpha_{j}) f_{0} + + 4\Delta_{j}^{2}f_{1} - Q_{1}^{2}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}) G_{1}^{(1)}(x,\eta,\alpha_{j}$$

Page 57.

$$\begin{array}{c} \times G_{1}^{(1)}(x,\ \eta,\ a_{j}) - \Delta_{j}^{2}G_{3}^{(2)}(x,\ \eta,\ a_{j})\} \frac{d\eta}{Vy-\eta} \\ L_{1,\ j}^{(1)}(x,\ y,\ a_{j}) = -\frac{\eta^{2}}{2\beta}\ V.\ p. \int\limits_{\chi(x)}^{g} \frac{\eta Q_{0}(\eta,\ a_{j})\ C(\eta,\ a_{j})}{\sqrt{y-\eta^{2}V}\ Z_{2}(x,\ \eta)} d\eta - \\ -\frac{1}{\beta} \int\limits_{\chi(x)}^{g} |[2\beta^{p}Q_{1}(x,\ \eta,\ a_{j}) - a_{j}^{2}\eta\ e] + 2(2\gamma-1)\ \Delta_{j}Z_{2}(x,\ \eta)] \times \\ \times \frac{C(\eta,\ a_{j})\ d\eta}{Vy-\eta} VZ_{3}(x,\ \eta) - \frac{2\beta^{2}a_{j}e}{3\Delta_{j}\gamma} [2(3\gamma-4)\ \Delta_{j}y + (4-3\gamma)[2\beta^{2}Q_{3}(x,\ a_{j}) - a_{j}^{2}\eta\ e] \int_{-1}^{g} |+ \\ +\frac{2}{\beta} \nabla a_{j}e \int\limits_{\chi(x)}^{g} \left\{ 2\beta^{3}\int_{-1}^{U_{1}}(p_{2}) - \frac{1}{2\gamma ea_{j}^{2}} [2\beta^{2}Q_{1}(x,\ \eta,\ a_{j}) - \\ -a_{j}^{2}\gamma e] G_{1}^{(1)}(x,\ \eta,\ a_{j}) \right\} \frac{d\eta}{Vy-\eta} - \frac{2\beta\Delta_{j}}{3a_{j}} \int\limits_{\chi(x)}^{g} \frac{G_{1}^{(2)}(x,\ \eta,\ a_{j})}{Vy-\eta} d\eta - \\ -a_{j}\nabla e \int_{1}^{g} -\frac{a_{j}\nabla e}{\beta} \int\limits_{\chi(x)}^{g} \left\{ \beta^{3}\int_{0} - \frac{1}{2}\ Q_{1}(x,\ \eta,\ 0)\ G_{2}^{(0)}(x,\ \eta,\ a_{j}) - \\ -\frac{\beta^{2}}{2}\ G_{2}^{(1)}(x,\ \eta,\ a_{j}) \right\} \frac{d\eta}{Vy-\eta} - \frac{2\beta\Delta_{j}}{Vy-\eta} \left\{ (35) \right\} \\ L_{1,\ j}^{(1)}(x,\ y,\ a_{j}) = -\frac{\nabla^{3}}{2\beta^{3}}\ V.\ p. \int\limits_{\chi(x)}^{g} \frac{\eta Q_{0}^{2}(\eta,\ a_{j})\ C(\eta,\ a_{j})d\eta}{Vy-\eta^{3}V} \frac{1}{Z_{1}(x,\ \eta)} - \\ -\frac{1}{\beta^{3}} \int\limits_{\chi(x)}^{g} [2(2\gamma-1)\Delta_{j}[2\beta^{2}\ Q_{1}(x,\ \eta,\ a_{j}) + \Delta_{j}Q_{1}(x,\ \eta,\ 0)]\ Z_{3}(x,\ \eta) + \\ +\beta^{3} \left[\beta^{2}Q_{1}^{2}(x,\ \eta,\ a_{j}) + 2\Delta_{j}Q_{1}(x,\ \eta,\ a_{j}) Q_{1}(x,\ \eta,\ 0)\right] + (4\gamma-1) \times \\ \times \Delta^{3}Z_{2}^{2}(x,\ \eta) \frac{1}{\sqrt{y-\eta}} \frac{C(\eta,\ a_{j})d\eta}{\sqrt{y-\eta}\sqrt{Z_{1}(x,\ \eta)}} - \frac{2a_{j}e}{15\gamma\Delta_{j}} \left\{3(15\beta^{4}+10\beta^{2}+1)\right\} \\ -2\beta^{3} \left\{\beta^{3}\left(\alpha_{j}^{2}(x,\ a_{j}) - 10\nabla a_{j}^{2}(3\nabla-2)\ e\right\} \left\{\Delta_{j}J_{1}+a_{j}^{2}eJ_{0}-12\beta^{2}(3\gamma-2)\ e\right\} \left\{\Delta_{j}J_{1}+a_{j}^{2}eJ_{0}-12\beta^{2}(3\gamma-2)\ e\right\} \left\{\Delta_{j}J_{1}+a_{j}^{2}eJ_{0}-12\beta^{2}(3\gamma-2)\ e\right\} \left\{\Delta_{j}J_{1}+a_{j}^{2}eJ_{0}-12\beta^{2}(3\gamma-2)\ e\right\} \left\{\Delta_{j}J_{1}+a_{j}^{2}eJ_{0}-12\beta^{2}(3\gamma-2)\ e\right\} \left\{\Delta_{j}J_{2}+\alpha_{j}J_{2}+\beta_{j}J_{0}+\alpha_{j}$$

Fage 58.

$$+\frac{4\eta}{\Delta_{f}} \left[ 5\eta \omega \theta_{f}^{2} - 6PQ_{h}(\omega, \omega_{f}) \right] Q_{h}(\omega, \omega_{f}) f_{-1} \left[ \frac{1}{4} \right] + \frac{\eta \omega_{f}}{\beta} \int_{I_{c}(x)}^{\beta} \left[ 4 \left[ 2P^{\mu}Q_{h}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}(x, \eta, \omega_{f}) \right] f_{-1}^{2}(\mu_{h}) + \frac{\eta \omega_{f}}{\beta} \int_{I_{c}(x)}^{\beta} \left[ 4 \left[ 2P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}(x, \eta, \omega_{f}) + \frac{\eta}{\gamma} \right] \frac{d\eta}{\sqrt{y - \eta}} \right] + \frac{1}{4} \int_{I_{c}(x)}^{\beta} \left[ 2P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}(x, \eta, \omega_{f}) \right] \frac{d\eta}{\sqrt{y - \eta}} \right] d\eta - \frac{2\Delta_{f}}{3\omega_{f}} \int_{I_{c}(x)}^{\beta} \left[ 2P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}(x, \eta, \omega_{f}) \right] \frac{d\eta}{\sqrt{y - \eta}} \right] d\eta - \frac{\omega_{f}\gamma \omega_{f}}{\sqrt{y - \eta}} \int_{I_{c}(x)}^{\beta} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \frac{\omega_{f}\gamma \omega_{f}}{\sqrt{y - \eta}} \right] d\eta - \frac{\omega_{f}\gamma \omega_{f}}{\sqrt{y - \eta}} \int_{I_{c}(x)}^{\beta} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}^{2}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}^{2}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}^{2}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}^{2}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}^{2}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) + \Delta_{f}Q_{h}^{2}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h}^{2}(x, \eta, \omega_{f}) \right] - \frac{1}{2} \left[ P^{\mu}Q_{h$$

Page 59.

$$-\frac{10\pi X}{(x)} - \frac{16\pi^{3}}{J_{1}} + \chi(x) [3\chi(x) - 5e] e f_{0} - \frac{1}{2\pi^{3}} V. p. \int_{\chi(x)}^{g} \sqrt{\frac{(e-\eta)^{3}\eta}{(y-\eta)^{3}} Z_{1}(x,\eta)} d\eta + \frac{15x^{2}}{16} \int_{\chi(x)}^{g} \frac{R_{2}(x,\eta) d\eta}{V \eta(y-\eta)} (38)$$

$$H_{1,1}(x,y) = -\frac{3}{49^{3}} \{5J_{3} - [3\chi(x) + 5e] J_{3} + 3e\chi(x) J_{1}\} - \frac{1}{29^{2}} V. p. \int_{\chi(x)}^{g} \sqrt{\frac{\eta(e-\eta)}{y-\eta}} \frac{d\eta}{V Z_{1}(x,\eta)} + \frac{9x}{4} \int_{\chi(x)}^{g} \sqrt{\frac{\eta}{y-\eta}} R_{2}(x,\eta) d\eta$$

$$H_{0,1}(x,y) = \frac{J_{1}}{2} - \frac{1}{2} V. p. \int_{\chi(x)}^{g} \sqrt{\frac{\eta^{3}(e-\eta)}{(y-\eta)^{3}} Z_{2}(x,\eta)} d\eta + \frac{3}{2} \int_{\chi(x)}^{g} \sqrt{\frac{\eta}{y-\eta}} R_{2}(x,\eta) d\eta$$

$$H_{0,2}(x,y) = \frac{1}{2} J_{3} - \frac{1}{2} V. p. \int_{\chi(x)}^{g} \sqrt{\frac{\eta^{3}(e-\eta)}{(y-\eta)^{3}} Z_{2}(x,\eta)} d\eta + \frac{5}{2} \int_{\chi(x)}^{g} \sqrt{\frac{\eta^{3}(e-\eta)}{y-\eta}} R_{2}(x,\eta) d\eta$$

$$N_{0}(x,y) = -J_{0} + 2J_{1} - \frac{\eta}{2} V. p. \int_{\chi(x)}^{g} \frac{V(e-\eta)\eta}{V(y-\eta)^{3}} \frac{d\eta}{Z_{2}(x,\eta)} + \frac{1}{49} \int_{\chi(x)}^{g} \frac{J_{0}(x,\eta) d\eta}{Vy-\eta} - \frac{1}{49} \int_{\chi(x)}^{g} \frac{J_{0}(x,\eta)}{Vy-\eta} \frac{d\eta}{Vy-\eta}$$

$$M_{1}(x,y) = \frac{1}{29^{3}} [e^{y}J_{0} - e(3\eta + 2) J_{1} + 4\eta J_{2}] - \frac{1}{29^{3}} [(3(\eta - 2)e^{-\eta}) V(e-\eta) \frac{1}{2} \frac{d\eta}{J_{1}} - \frac{1}{29^{3}} [(3(\eta - 2)e^{-\eta}) V(e-\eta) \frac{1}{2} \frac{d\eta}{J_{2}} - \frac{1}{29^{3}} [(3(\eta - 2)e^{-\eta}) V(e-\eta) \frac{1}{2} \frac{1}{29^{3}} [(3(\eta - 2)e^{-\eta}) V(e-\eta) \frac{1}{2} \frac{1}{29^{3}} [(3(\eta - 2)e^{-\eta}) V(e-\eta) \frac{1}{2}$$

DQC = 78184200

Page 60.

$$-8\beta^{2}\eta I_{0} + 8\beta^{2}I_{1} \frac{d\eta}{Vy-\eta} - \frac{1}{4\beta} \int_{\chi(x)}^{\eta} \{[4\beta^{2}(\eta-x)^{2} - 3(\nabla-2)e(\eta-x) - 2e^{2}]G_{2}^{(0)}(x,\eta) + [8\beta^{2}(\eta-x) - 3(\nabla-2)e]G_{2}^{(1)}(x,\eta) + 4\beta^{2}G_{2}^{(1)}(x,\eta) \} \frac{d\eta}{Vy-\eta}$$

$$-3(\nabla-2)e]G_{2}^{(1)}(x,\eta) + 4\beta^{2}G_{2}^{(1)}(x,\eta) \} \frac{d\eta}{Vy-\eta}$$

$$N_{2}(x,y) = -\frac{1}{2\beta^{4}} \{e^{2}J_{0} - 2(3\nabla+1)e^{2}J_{1} + \nabla(5\nabla+8)eJ_{2} - 6\nabla^{2}J_{3}\} - \frac{\nabla}{2\beta^{4}} V. p. \int_{\chi(x)}^{y} \frac{(\nabla\eta-e)^{2}V}{Vy-\eta^{3}V} \frac{\eta(e-\eta)}{Z_{2}(x,\eta)} d\eta - \frac{1}{\beta} \int_{\chi(x)}^{y} \{[2e^{2} + 5(\nabla-2)e\eta - 9\beta^{2}\eta^{2}]I_{0} + [18\beta^{2}\eta - 5(\nabla-2)e]I_{1} - 9\beta^{2}I_{2}\} \frac{d\eta}{Vy-\eta} - \frac{1}{4\beta} \int_{\chi(x)}^{y} \{[6\beta^{2}(\eta-x)^{3} - 5(\nabla-2)e(\eta-x)^{2} - 4e^{2}(\eta-x)]G_{2}^{(0)}(x,\eta) + 2[9\beta^{2}(\eta-x)^{2} - 5(\nabla-2)e(\eta-x) - 2e^{2}]G_{2}^{(1)}(x,\eta) + + 16\beta^{2}(\eta-x) - 5(\nabla-2)e]G_{2}^{(2)}(x,\eta) + 6\beta^{2}G_{2}^{(2)}(x,\eta) \frac{d\eta}{Vy-\eta}$$

Page 61.

In formulas (17)-(44) are introduced the following designations:

$$f_{k} = \int_{\chi(x)}^{x} \frac{\eta^{k} d\eta}{V(y-\eta) \, \eta(e-\eta)(\eta+\beta^{2}x-e)}$$

$$f_{-1} = \int_{\chi(x)}^{y} \frac{d\eta}{\left(-\frac{\alpha_{j}^{2}e}{\Delta_{j}}-\eta\right) V(y-\eta) \, \eta(e-\eta)(\eta+\beta^{2}x-e)}$$

$$G_{1}^{(m)}(x, \, \eta_{i}, \, \alpha_{j}) = \frac{2\alpha_{j}^{2}\nabla e}{\nabla_{j}} \int_{-2}^{(m)}(p_{3}) - (\alpha_{j}^{2}+\beta^{2}) \int_{-1}^{(m)}(p_{3}) + \alpha_{j}^{2} \int_{-1}^{(m)}(p_{3}) + \beta^{2} \int_{-1}^{(m)}(p_{3}) + \beta^{2$$

$$p_1 = \eta - e, \quad p_2 = \frac{e}{\beta^n} + \eta, \quad p_3 = \frac{Q_1(0, \eta, \alpha_j)}{\Delta_j}$$

$$I_m = \int_{\tilde{X}(\eta)}^{\varepsilon} \frac{\xi^m d\xi}{\sqrt{(x - \xi)(e + \xi - \eta)\left(\frac{e}{\beta^2} + \eta - \xi\right)}}$$

$$I_m^{(m)}(\rho) = \int_{\tilde{X}(\eta)}^{\varepsilon} \frac{(x - \xi)^m d\xi}{(\rho - \xi)^n \sqrt{(x - \xi)(e + \xi - \eta)\left(\frac{e}{\beta^2} + \eta - \xi\right)}}$$

In the last/latter integral p, are taken values  $p_1$ ,  $p_2$  and  $p_3$ .

All integrals of types  $l_m$ ,  $l_m^{(m)}$ ,  $l_k$  and  $l_{-1}$  are expressed as elliptical integrals

$$I_{0} = \frac{2\beta}{V \nabla e} \mathbb{E}(\delta, q)$$

$$I_{1} = \frac{2}{\beta V \nabla e} [-Q_{1}(x, \eta, 0) \Pi(\delta, q^{2}, q) + Q_{1}(0, \eta, 0) \mathbb{E}(\delta, q)]$$

$$I_{2} = -\frac{2\nabla}{3\beta^{3}} V \frac{1}{\eta(e-\eta)} \frac{1}{Z_{2}(x, \eta)} - \frac{4Q_{1}(x, \eta, 0)}{3\beta^{3} V \nabla e} [e(1-\beta^{2}) + \beta^{2}x + 2\beta^{2}\eta] \Pi(\delta, q^{2}, q) + \frac{2}{3\beta^{3} V \nabla e} [2Q_{1}^{2}(0, \eta, 0) + \beta^{2}(x + \eta - - e) Q_{1}(0, \eta, 0) + \beta^{4}(e-\eta) x] \mathbb{E}(\delta, q)$$

$$I_{-1}^{(0)}(p_{2}) = -\frac{2\beta \Delta_{j}^{2}}{\alpha_{j}^{2} V \nabla e^{3}} \frac{1}{Q_{1}(x, \eta, \alpha_{j})} \Pi(\delta, \frac{\alpha_{j}^{2} e \nabla q^{2}}{\beta^{3} Q_{1}(x, \eta, \alpha_{j})}, q) + \frac{2\beta^{3}\Delta_{j}}{\alpha_{j}^{2} V \nabla e} \mathbb{E}(\delta, q)$$

$$I_{-1}^{(0)}(p_{2}) = -\frac{1}{\alpha(x, \eta, \alpha_{j})} \left[ \frac{Q_{1}(x, \eta, 0)}{V \nabla} \Pi(\delta, q^{2}, q) + \frac{\alpha_{j}^{2} V \nabla}{\beta \Delta_{j}} \mathbb{E}(\delta, q) + \frac{2\beta^{3}\Delta_{j}}{\alpha_{j}^{2} V \nabla^{2}Q_{1}(x, \eta, \alpha_{j})} \Pi(\delta, \frac{\alpha_{j}^{2} \nabla e q^{2}}{\beta^{2}Q_{1}(x, \eta, \alpha_{j})}, q) - \frac{\beta^{3}\Delta_{j}}{\alpha_{j}^{2} V \nabla^{2}} b(x, \eta, \alpha_{j}) \mathbb{E}(\delta, q) + \frac{\Delta_{j} V \eta(e-\eta) Z_{2}(x, \eta)}{\beta Q_{0}(\eta, \alpha_{j})} \right]$$

$$J_{0} = \frac{2}{\beta V x y} \mathbb{E}\left(\frac{\pi}{2}, q_{1}\right)$$

$$J_{1} = \frac{2}{\beta V x y} \left\{ (y - e) \Pi\left(\frac{\pi}{2}, x, q_{1}\right) + e \mathbb{E}\left(\frac{\pi}{2}, q_{1}\right) \right\}$$

$$J_{1} = \frac{2}{\beta V x y} \left\{ \chi(x)(e-y) \Pi\left(\frac{\pi}{2}, \frac{ex}{3}, q_{1}\right) + (Z_{2} + e)(e-y) \times \frac{e^{x}}{\beta^{2}} \right\}$$

DOC = 78184200

Page 62.

$$\times \Pi\left(\frac{\pi}{2}, x_{i}, q_{i}\right) + \left[\beta^{2}x(e-y) - 2e^{x}\right] \mathbb{E}\left(\frac{\pi}{2}, q_{i}\right)$$

$$J_{0} = -\frac{1}{2} \frac{ey \chi(x)}{\beta \sqrt{xy}} \mathbb{E}\left(\frac{\pi}{2}, q_{i}\right) - \frac{3\chi(x)}{4\beta \sqrt{xy}} (Z_{i} + e) \left[(e-y) \times \right.$$

$$\times \Pi\left(\frac{\pi}{2}, \frac{ex}{y}, q_{i}\right) + y \mathbb{E}\left(\frac{\pi}{2}, q_{i}\right) \right]$$

$$J_{-1} = \frac{2\Delta_{j}(e-y)}{\beta^{2}e Q_{0}(y, x_{j})} \Pi\left(\frac{\pi}{2}, \frac{e\beta^{2}x}{Q_{0}(y, x_{j})}, q_{i}\right) + \frac{2}{\beta^{2}e} \mathbb{E}\left(\frac{\pi}{2}, q_{i}\right)$$

where they are designated

$$\delta = \arcsin \sqrt{\frac{e(\eta + \beta^2 x - e)}{\beta^2 \eta(x - \eta + e)}}, \quad q = \sqrt{\frac{\beta^2 (x - \eta + e)}{\gamma e}}$$

$$x = \frac{y + \beta^2 x - e}{\beta^2 x}, \quad q_1 = \sqrt{\frac{e(y + \beta^2 x - e)}{\beta^2 x y}}$$

$$a(x, \eta, \alpha_j) = \frac{1}{\beta^2} \left(\beta^2 p_3^2(\eta, \alpha_j) - [Q_1(0, \eta, 0) + \beta^2(x + \eta - e)] p_3^2(\eta, \alpha_j) + [Q_1(0, \eta, 0)(x + \eta - e) - \beta^2(e - \eta) x] p_3(\eta, \alpha_j) + (e - \eta) Q_1(0, \eta, 0) x\}$$

$$b(x, \eta, \alpha_j) = \frac{1}{\beta^2} \left\{3\beta^2 p_3^2(\eta, \alpha_j) - 2[Q_1(0, \eta, 0) + \beta^2(x + \eta - e)] p_3(\eta, \alpha_j) + Q_1(0, \eta, 0)(x + \eta - e) - \beta^2(e - \eta) x\right\}$$

Through E and P are designated the elliptical integrals respectively of the first and third kinds.

Page 63.

Por a region o, (Fig. 1) the formula of the coefficients rotary

DQC = 78184200

PIGE 2524

Taking this opportunity, I express my sincere appreciation to Frof. R. A. Mezhlumyan for large interest in task and valuable observations, and also to Dzh. A. Prutyunyan for aid during checking of the obtained results.

Scientific-Research Institute of Automation

Kirovakan

Received 1 II 1971

Page 64.

## NON-STATIONARY DOWNWASH BEHIND A TRIANGULAR WING IN SUPERSONIC MOTION

R. Sh. SOLOMONIAN

Summary

Rated formulae for non-stationary downwash behind a thin slightly curved wing which has a traingular formula in the plan and supersonic edges whose apex is turned forward when it moves in the ideal compressible fluid with supersonic speed are given in this paper.

These formulae are obtained for area points situated on the plane of the wing between the disturbance waves, their reflection from the back edge and the back edge itself at small Strouhal numbers.

The above formulae make it possible to continue the downstream calculations at infinitum according-to the formulae previously derived for this part of the wing plane.

#### FEFERENCES

- 1. S. M. Belotserkovskiy. Three-dimensional/space unsteady motion of lifting surfaces. PMM, Vcl. XlX, iss. 4, 1955.
- (2). R. A. Mezhlumyan, F. Sh. Sclomonyan. Method of determining the unsteady wing downwash of final spread/scope during supersonic motion. Izv. AS ARM SSR, the series of mechanic, Vol. XXIII, No 6, 1970.
- (3). N. N. Kislyagin. Ccefficients rotary derivative tapers, created in flow by wing, during unsteady motion. Izv. of the AS USSR, OTN,

DOC = 78184200

mech. and machine-building, No 4, 1961.

- (4). "General theory of high-speed aerodynamics". Series "High-speed aerodynamics and jet-propulsion technology". Translated from Engl.

  IL, M., 1959.
- (5). "Contemporary state of high-speed aerodynamics", General editor:
  L. Howard, IIL, 1955.
- (6). I. S. Gradshteyn and I. M. Ryzhik. Tables of integrals, sums, series and products. Fizzatçiz, M., 1963.

#### DISTRIBUTION LIST

### DISTRIBUTION DIRECT TO RECIPIENT

ORGANIZATION		MICROFICHE	ORGANIZATION		MICROFICHE
A205	DMATC	1	E053	AF/INAKA	1
A210	DMAAC	2	E017	AF/RDXTR-W	1
B344	DIA/RDS-3C	9	E403	AFSC/INA	1
C043	USAMIIA	1	E404	AEDC	1
C509	BALLISTIC RES LABS	1	E408		1
C510	AIR MOBILITY R&D	1	E410	ADTC	1
	LAB/FIO				
	PICATINNY ARSENAL	1		FTD	
C535	AVIATION SYS COMD	1		CCN	1
C591	FSTC	5		ASD/FTD/NII	S 3
C619	MIA REDSTONE	1		NIA/PHS	1
D008	NISC	1		NIIS	2
H300	USAICE (USAREUR)	1			
P005	DOE	1			
P050	CIA/CRB/ADD/SD	1			
NAVORDSTA (50L)		1			
NASA/KSI		1			
AFIT/LD		1			
LLL/Code L-389		ī			